




Research Article

Running out of empty space: environmental lidar and the crowded ancient landscape of Campeche, Mexico

Luke Auld-Thomas^{1,2,*} , Marcello A. Canuto^{1,3}, Adriana Velázquez Morlet⁴, Francisco Estrada-Belli³, David Chatelain¹, Diego Matadamas¹, Michelle Pigott¹ & Juan Carlos Fernández Díaz⁵


¹ Department of Anthropology, Tulane University, New Orleans, USA


² Department of Anthropology, Northern Arizona University, Flagstaff, USA

³ Middle American Research Institute, Tulane University, New Orleans, USA

⁴ Instituto Nacional de Antropología e Historia (INAH), Ciudad de México, México

⁵ National Center for Airborne Laser Mapping (NCALM), University of Houston, USA

* Author for correspondence  lauldth@tulane.edu



As airborne lidar surveys reveal a growing sample of urbanised tropical landscapes, questions linger about the sampling bias of such research leading to inflated estimates of urban extent and population magnitude. ‘Found’ datasets from remote sensing conducted for non-archaeological purposes and thus not subject to archaeological site bias, provide an opportunity to address these concerns through pseudorandom sampling. Here, the authors present their analysis of an environmental lidar dataset from Campeche, Mexico, which reveals previously unrecorded urbanism and dense regional-scale settlement. Both characteristics, the authors argue, are therefore demonstrably ubiquitous across the central Maya Lowlands.

Keywords: Central America, Maya Lowlands, remote sensing, lidar, low-density urbanism, settlement patterns

Introduction

Archaeologists increasingly recognise that the global tropics and sub-tropics were home to a rich variety of urban settlement traditions in antiquity. While many of these settlement systems shared a tendency toward spatial dispersion—what is frequently called ‘low-density urbanism’ (Fletcher 1995, 2009, 2012)—there is a burgeoning recognition that the ancient world’s low-density urban landscapes were themselves heterogeneous, with substantial gradients in

Received: 24 August 2023; Revised: 6 February 2024; Accepted: 14 March 2024

© The Author(s), 2024. Published by Cambridge University Press on behalf of Antiquity Publications Ltd. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.

settlement density within settlements, between settlements and their hinterlands, and across subregions (Chase *et al.* 2011; Evans *et al.* 2013; Fisher *et al.* 2017; Canuto *et al.* 2018; Jackmond *et al.* 2018; Prümers *et al.* 2022; Thompson 2023; Rostain *et al.* 2024). Concurrently, archaeologists in many of these same regions are confronting ‘dense pasts’ (see Evans *et al.* 2023), where the accumulation of research has simply turned up more of everything than the discipline contemplated previously: more settlements, more cities, more landesque capital.

In recent years these two developments in the archaeology of tropical urbanism—the recognition of high-amplitude variability on the one hand, and of a dense past on the other—have come into tension with one another, such that where some researchers see evidence for dense regional-scale populations inhabiting thoroughly urbanised landscapes, others see evidence of a biased sample that over-represents one end of the settlement density gradient. It thus remains an open question whether those areas as yet unmapped by archaeologists hold further evidence of a dense past or reveal a less densely occupied corrective.

Here we seek to answer this question for the central Maya Lowlands (see Canuto *et al.* 2018: fig. 1), one of the ancient world’s best-known, but also most contentious, examples of tropical urbanism. Our aim in this article is to assess whether overall settlement density and local-scale variability in settlement density (i.e. urbanisation) have been over-estimated by archaeologically motivated, ‘site focused’ surveys. To do this we enlist a new expedient and pseudorandom regional-scale sample of settlement drawn from remote sensing data across one of the Maya Lowlands’ largest ‘blank spots’: eastern Campeche, Mexico.

Since at least the 1940s, researchers have known that the Classic period Maya (AD 250–900) transformed the rugged interior of the Mexican state of Campeche into a densely settled and extensively engineered landscape (Merwin 1913; Ruppert & Denison 1943). A series of projects undertaken near the modern town of Xpuhil since the 1970s demonstrate that this region was both crowded with settlement and heavily engineered through durable landscape modifications such as terraces and field walls (Eaton 1975; Turner II 1978; Thomas 1981; Suárez Aguilar 2007; Arnauld *et al.* 2013; Lemonnier & Vannière 2013). Nevertheless, for most of the twentieth century, archaeological research in east-central Campeche and adjacent Quintana Roo was scant compared to other parts of the Maya Lowlands, such that the authoritative database of known and geolocated sites (Witschey & Brown 2014) shows large swaths of the region to be essentially *terra incognita*.

In the past decade a growing number of field research projects in Campeche have employed airborne lidar survey (e.g. Reese-Taylor *et al.* 2016; Hernández Gómez 2021; Šprajc *et al.* 2021), which has emerged as a powerful tool for settlement archaeology, capable of documenting entire landscapes in minute detail even under dense forest cover (e.g. Chase *et al.* 2011; Canuto *et al.* 2018). Separately, researchers have also examined lidar datasets collected for non-archaeological purposes, notably those for ecological monitoring (Schroder *et al.* 2020; Hutson *et al.* 2021) and civil engineering (Instituto Nacional de Antropología e Historia 2020; Pérez Rivas *et al.* 2023). Collectively referred to as environmental lidar, these expedient datasets have the advantage of being effectively random with regards to the archaeological record; as the data were collected without regard for archaeological questions they are unlikely to contain the same biases as archaeologically motivated surveys. In the Maya Lowlands and elsewhere (Golden *et al.* 2016; García Sánchez 2018; Schroder *et al.* 2020; Kokalj & Mast 2021; Peripato *et al.* 2023; Stauffer *et al.* 2023), archaeological

re-analysis of these environmental lidar datasets represents a promising new development in regional sampling methodology (Plog & Flannery 1976; see also Banning 2021) with an attractive combination of high spatial resolution, broad spatial coverage, limited archaeological bias and low (or no) cost. In Mexico, some environmental lidar datasets have been made freely available to the public—notably the transects from NASA’s G-LiHT mission (Goddard’s Lidar, Hyperspectral and Thermal Imager mission) (Cook *et al.* 2013)—in addition to the government’s own high-resolution lidar data, fostering collaboration among researchers as well as open science, more generally.

Here, we present an archaeological analysis of approximately 122km² of high-quality airborne lidar data collected in 2013 as part of a forest monitoring project called Alianza M-REDD+ (Reducing Emissions from Deforestation and Degradation in Mexico), spearheaded by the Nature Conservancy, Mexico. This dataset—hereafter ‘Alianza’—uniquely comprises both transect and areal (quadrat or block) survey samples, permitting a pseudorandom sample of settlement patterns in Campeche that is both longitudinal and contextual (Figure 1).

In this article, we introduce the dataset and provide an overview of the archaeological landscape that it reveals. We argue that the Alianza data demonstrate a range of ancient settlement densities comparable to those documented by site-focused archaeological survey in the region, from nearly vacant rural landscapes to dense urban areas. Despite this survey’s lack of archaeological “site bias” (c.f. Ringle *et al.* 2021: 46–47), we conclude that the Alianza datasets support the notion of a populous and urban ancient Maya landscape consistent with other datasets (Canuto *et al.* 2018; Schroder *et al.* 2020; Canuto & Auld-Thomas 2021, 2024). In other words, we suggest that dense regional-scale settlement and high-amplitude variability in local-scale settlement density (reflective of urbanisation) are characteristic of the central Maya Lowlands and not the product of sampling bias (contra Webster 2018). More detailed quantitative analysis of the archaeological features within this and other pseudorandom datasets will be published elsewhere in the future.

The dataset

The lidar data were collected in 2013 by the Mexican firm CartoData using a Riegl LMS-Q780 sensor, processed by the Woods Hole Research Center (WHRC) and made freely available by Alianza M-REDD+ (Walker 2015). The dataset consists of three transects and three survey blocks (Figure 1). The transects are roughly 275m wide and approximately 213km long, for a total coverage area of 58.3km². The survey blocks total 64.1km² (Table 1) and were conducted in three areas: 1) south of the town of Xpuhil, near the well-known site of Río Bec; 2) adjacent to the villages of Dos Lagunas and Bel Ha where several patio groups have been reported along the highway (Merk 2004); and 3) near the town of Ucum in northernmost Campeche, an area with no reported sites. A fourth survey block, measuring about 174km², falls in the Puuc-Bolonchen hills of the state of Yucatan (see Ringle *et al.* 2021), and is not considered further here.

Analytical methods

In the original Alianza digital terrain models, larger archaeological mounds were vertically ‘truncated’, indicating that a subset of real ground points in the lidar point cloud had not




Figure 1. Alianza lidar survey areas in Campeche (figure by authors).

been classified as such. We reprocessed the original point clouds using parameters developed at the National Center for Airborne Laser Mapping (NCALM) specifically for archaeological prospection (Fernandez-Diaz *et al.* 2014), producing a digital terrain model with a spatial resolution of 1m per pixel (ground-point density: 5.01points/m²). This dramatically improved the representation of ancient architecture, though the overall visibility of archaeological features across the dataset was only slightly improved. Digital terrain models were

Table 1. Settlement data in Alianza lidar in Campeche. Density is measured in structures per square kilometre.

Survey unit	Area (km ²)	Structures	Density (strs/km ²)
Block 1	13.77	744	54.0
Block 2	16.66	1732	104.0
Block 3	33.63	667	19.8
Transect 1	30.31	1937	63.9
Transect 2	14.62	1038	71.0
Transect 3	13.39	646	48.3
Total	122.37	6764	55.3

visualised primarily using Simple Red Relief (Auld-Thomas 2022), which highlights slopes in red and local elevation differences in a gradient of dark blue-green to bright yellow (e.g. Figures 4 & 7), and the Visualization for Archaeological Topography (Verbovšek *et al.* 2019), which blends a traditional computational hillshade with additional terrain parameters to enhance subtle topographic details. All features of archaeological interest were manually digitised as vector polygons in QGIS and ArcGIS Pro. Digitized features were reviewed by at least three analysts working independently; where analysts disagreed, features were reevaluated by all parties and then included or excluded from the final dataset by majority vote.

To our knowledge, archaeological fieldwork has been conducted only in the Puuc block and not any of the other areas intersected by the survey; and no field validation of the analysis presented here has yet taken place. This means that our analysis carries three caveats. First, over the decade since lidar was first deployed in the service of Mesoamerican archaeology, in no instance has field validation resulted in a net decrease in the total number of buildings identified (e.g. Reese-Taylor *et al.* 2016; Canuto & Auld-Thomas 2020; Thompson 2020; Garrison *et al.* 2022), because false positives are reliably offset by a greater number of false negatives—small, unrounded or barely rounded buildings that are seldom detectable in digital terrain models. Nor in that time has field validation of lidar-derived settlement data resulted in a meaningful alteration to the observed spatial distribution of (mounded, surface-visible) settlement.

The second caveat concerns the long-recognised prevalence of rockpiles (or ‘chich’ mounds) throughout the Alianza coverage area. Although their exact use is unknown (Turner II 1983; Dunning 1992; Kepecs & Boucher 1996; Lemonnier & Vannièrè 2013; Hutson *et al.* 2021), most of these rockpiles do not appear to have fulfilled a residential function. Even so, their general shape (as depicted in lidar data) is similar to that of small house-mounds. Despite efforts to mitigate the impact of this ambiguity during our digitisation process, our identification efforts likely mistook some rockpiles for structures, leading to marginally inflated structure tallies overall.

The net result of these first two caveats is that all feature counts and related analyses of density presented here should be regarded as preliminary values that will inevitably be revised through subsequent analyses, both in the field and the computer lab. Overall feature counts will almost certainly increase, but the proportion of features classified as buildings or some other anthropogenic feature, such as rockpiles, is also likely to change. We maintain, however, that the general characteristics and spatial patterning of settlement can be reliably described from our initial remote analysis.

The final caveat concerns chronology and contemporaneity. In the central lowlands, archaeologists have long recognised a “tremendous Late Classic bias” (Rice & Puleston 1981: 137) in buildings visible on the surface; nearly every archaeological project in the region has reported that between 75 and 95 per cent of excavated buildings have Late Classic components (e.g. Canuto & Auld-Thomas 2024). Similar chronological summaries are not available for the northern lowlands, although the chronological picture there is generally regarded as more complex and there is far more Terminal Classic and Postclassic occupation compared with the central lowlands. The lack of chronological control in the area sampled by the Alianza survey means that the northern and southern portions of the dataset should not be considered synchronic, though elements of each surely overlap in time.

Settlement patterns

General observations

Block 1, located just south and east of the important site of Río Bec, includes one probable example of the distinctive paired pseudopyramid architectural style that takes its name from that site (Figure 2a). The character of settlement, including the ubiquity of agricultural infrastructure, is identical to that recorded by pedestrian survey at Río Bec: a sprawling ‘dense




Figure 2. Left) Block 1 survey block, with building footprints in black and platform footprints in purple; top right) probable Río Bec-style complex marked by *; bottom right) sinkhole with associated quatrefoil ditch (figure by authors).

rural' agro-settlement landscape with little of the imposing public architecture so common elsewhere in the Maya region (Arnauld *et al.* 2013; Lemonnier & Vanni re 2013). One notable exception, however, is an architectural complex adjacent to a deep sinkhole giving access to a partially collapsed cave system. The architectural complex is encircled by a narrow ditch, the small scale of which, together with the lack of an associated embankment, precludes a defensive interpretation (Figure 2b). The ditch is in the shape of a quatrefoil, canonically the shape used to represent caves and sinkholes in Mesoamerican art (Guernsey 2010). This unusual pairing of prominent sinkhole and quatrefoil ditch warrants future investigation.

Block 2 directly overlies the epicentre of a major urban area that was previously unknown to the scientific community. This city, which we call Valeriana (Figures 3, 4 & 5) after an adjacent freshwater lagoon, encompasses two major hubs of monumental architecture 2km apart, linked by continuous dense settlement and landscape engineering. The discovery of Valeriana highlights the fact that there are still major gaps in our knowledge of the existence or absence of large sites within as-yet unmapped areas of the Maya Lowlands. Settlement and agricultural infrastructure completely fill the 16.6km² survey area. The style and density of architecture closely matches that seen in the Chactun-Tamchen area, some 20km to the south-west ( prajc *et al.* 2021), as well




Figure 3. Block 2, showing the major site of Valeriana. Structure footprints in black, platform footprints in purple. Two mound groups adjacent to the highway were reported by Merk (2004) (figure by authors).




Figure 4. Detail of Valeriana site core, in the north-east corner of Block 2 (figure by authors).

as that recently recorded by archaeological salvage work on the Maya Train project (Pérez Rivas, *pers. comm.*).

The larger of Valeriana's two monumental precincts has all the hallmarks of a Classic Maya political capital: multiple enclosed plazas connected by a broad causeway; temple pyramids; a ballcourt; a reservoir formed by damming an arroyo (a seasonal watercourse); and a probable E-Group assemblage, an architectural arrangement that generally indicates a founding date prior to AD 150 (Freidel *et al.* 2017). Furthermore, the same style of curved, 'amphitheatre' residential patios first identified by Ruppert and Denison (1943) at the site of Pechal and reported in the northern sector of Šprajc and colleagues' (2021) lidar survey are also present at Valeriana, making this a strongly characteristic regional style. Valeriana's architectural epicentre occupies the north-eastern extreme of Block 2, indicating that the settled area associated with it continues beyond the margins of the Alianza dataset. It likely continues at least 5km north, where Transect 2 records identical density and style of buildings and agricultural infrastructure (see below).

Block 3, by contrast, has sparse and modest settlement (Figure 6), consisting of scattered or loosely clustered residences with no monumental architecture and limited investment in water storage. Notably, the settlement here does include so-called 'annular structures', ring-shaped structures that are common in the Puuc region and were used in the production of lime plaster (Seligson *et al.* 2019).




Figure 5. Settlement and agricultural land modification in Block 2, visualisation for archaeological topography. Compare with figures in Šprajc et al. 2021 (figure by authors).

Settlement density

With a total of 6764 structures (strs) in the entire dataset, the aggregate settlement density is $55.3\text{strs}/\text{km}^2$, comparable to the $49\text{strs}/\text{km}^2$ recently reported by Schroder and colleagues (2020) for a portion of the G-LiHT transects in the same region. These figures are relevant to assessments of regional-scale settlement density and population history—for instance, they exceed regional-scale settlement density reported for any part of Belize or Guatemala (Canuto & Auld-Thomas 2024). However, they explain little about the degree to which a settlement system was urbanised, which requires an appreciation of local-scale variability or density gradients. To accommodate the distinct spatial qualities of quadrat and transect data, we assess density gradients separately for the Alianza blocks and transects.

To calculate settlement density gradients in the survey blocks, we used a kernel density estimation, fitting a quartic kernel with a 565m bandwidth over each structure point with an output cell size of 5m. We did the same for the only pedestrian surveys conducted at comparable scale and completeness in the same region, namely at Calakmul (May Hau et al. 1990), Oxpemul (Barnes et al. 2009) and Becan (Thomas 1981). Comparing these ‘site-centred’ block surveys with ‘random’ block surveys illustrates that the latter fit comfortably within the range of densities suggested by the former (Figure 7). Block 1 (range: $0\text{--}195\text{strs}/\text{km}^2$) falls between densities recorded at Oxpemul (range: $0\text{--}149\text{strs}/\text{km}^2$) and Becan (range: $0\text{--}264\text{strs}/\text{km}^2$),




Figure 6. Settlement in Block 3 (building footprints in black). Annular structures are associated with lime plaster production (Ringle et al. 2021). Details A, B and C illustrate Northern Lowland settlement morphology, compare with Figure 7 (figure by authors).

while Block 2 (range: 0–426strs/km²) displays a density gradient only surpassed by the behemoth site of Calakmul (range: 0–770strs/km²). Notably, the placement of Block 2 vis-à-vis Valeriana’s dense epicentre creates an edge effect precisely in the area where settlement density is greatest; that city’s urban core is undoubtedly both larger and denser than our kernel density estimation suggests. Block 3 (range: 0–84strs/km²) shows the lowest, most ‘rural’, settlement distribution of all six survey blocks; block-level density here is comparable to that reported for rural areas such as the Corona-Achiotal region in northern Guatemala (Canuto & Auld-Thomas 2021), though it is also more evenly distributed. In sum, Alianza’s three pseudorandom survey blocks recorded: 1) a low-density rural area; 2) a moderately dense periurban or ‘dense rural’ area; and 3) a large, dense city—precisely the range of variability suggested by ‘site-centred’ pedestrian and lidar surveys.

Due to the narrowness of the transects (between 200m and 300m), we devised a binning/segmentation procedure to assess variability in settlement density. Schroder and colleagues (2020) used a similar strategy in their analysis of the G-LiHT data, dividing the transects into roughly 2km² segments (approximately 8km in length). We reasoned that bins of that length would be unlikely to capture significant short-scale variation in settlement density, since even the largest Maya cities’ epicentres seldom exceed 4km in any linear dimension




Figure 7. Comparison of settlement density between six areas in central Campeche (figure by authors).

and are often bounded by vacant wetlands and sparsely occupied rugged terrain (Canuto *et al.* 2018); the effect of segments this long is to reduce a city’s analytical visibility. Consequently, we employ a segment size of 25ha (approximately 1km in length). These reveal greater variability in settlement density, allowing us to identify areas of nucleation (potential loci of ‘urbanising processes’) as well as areas with little or no settlement (‘hinterlands’).

Dense settlement is most broadly sustained at the eastern end of Transect 2, only 5km to the north of Block 2 and the site of Valeriana, suggesting that the region encompassing Block 2 and the eastern end of Transect 2 is part of an extended high-density settlement system (Figure 8C), like the one recorded by Šprajc and colleagues (2021) around Chactun—indeed, the Alianza and Chactun survey areas are close enough that they may simply capture parts of a single sprawling conurbation.

There are other areas where density appears to spike to levels typical of ‘urban’ landscapes. Two are found along Transect 1, at points near the known centres of Dzibilnocac and (somewhat farther north) Santa Rosa Xtampak (Figure 8A). A third localised spike in density corresponds to a previously unreported monumental complex at the extreme southern terminus of Transect 1 (Figure 8B); because this complex occupies the southern end of the transect, it is impossible to say how large it may be or how urban the settlement associated




Figure 8. Close-ups of transects; A) Transect 1; B) southern tip of Transect 1; C) eastern end of Transect 2. All densities expressed as structures per square kilometre (figure by authors).

with it was. Transect 3 records the lowest density of all three transects; nowhere does it intersect densities typical of urban areas, and its eastern leg has the longest stretch of low-density segments (range: 0–0.84strs/ha or 0–84strs/km²) in the entire dataset, matching the low-density Block 3 and suggesting the existence of a large and sparse settlement zone in north-easternmost Campeche.

Regional variability

The longitudinal nature of the three transects helps to reveal changes in settlement morphology, especially moving south to north (Figure 9). Throughout the southern half of the survey area,




Figure 9. Changes in settlement morphology from south to north. A) 18°57'0"N, 89°18'36"W, Transect 2, Central Hills; B) 19°1'48"N, 89°33'0"W, Transect 2, Central Hills; C) 19°10'48"N, 89°33'36"W, Transect 3, Central Hills; D) 19°32'24"N, 89°43'12"W, Transect 1, Chenes; E) 19°34'48"N, 89°42'36"W, Transect 1, Chenes (note high frequency of chich mounds); F) 20°0'0"N, 89°39'0"W, Transect 1, Puuc-Bolonchen Hills (figure by authors).

architectural remains strongly favour upland locations and generally consist of mounded structures arranged orthogonally or in a continuous arc around a built-up or levelled patio.

The same landform-crowding phenomenon documented by Canuto and Auld-Thomas (2021) in Guatemala is apparent in the Alianza survey blocks, with urban Block 2 having a higher settlement rate on marginal landforms (42strs/km²) than in the more rural Block 1 (10strs/km²) or sparsely settled Block 3 (8strs/km²). Agricultural infrastructure, in the form of field walls and terraces, is ubiquitous and occurs in close association with settlement. Wetland fields are notably absent from the Alianza survey data, although complexes of narrow drainage canals articulating with field walls are known from the adjacent Chactun region (Dunning *et al.* 2019; Šprajc *et al.* 2021).

Both the morphology of settlement remains and their topographic patterning change gradually but markedly above the 19th parallel (Turner II 1983), beginning in the northern part of what Dunning and colleagues (1998) call the Central Hills adaptive region. Terracing and field walls become less common, then all but disappear. Vernacular architecture changes as well: mounded residential patio groups are mostly replaced by large basal platforms greater than 200m² interspersed with less orthogonally arranged individual buildings. Both this architectural tradition and the absence of agricultural terracing are long-noted characteristics of settlement in the Puuc region and adjacent parts of the northern Maya Lowlands (Ringle *et al.* 2021), although field walls are common further north (e.g. Hutson 2016). The strong correlation of upland landforms and archaeological settlement also progressively breaks down north of the 19th parallel and is largely inverted north of Hopelchen; in the northern reaches of Transect 1, most buildings are in valley bottoms and on spurs or foot slopes—long noted as a central characteristic of settlement in the Puuc region (Dunning & Kowalski 1994).

Given the generally synchronous changes in architectural practice, settlement morphology, agricultural engineering and landform preference, we suggest that the northern limits of the Central Hills region—around the towns of Xkanha, Kancabchen and Ucum—is a reasonable benchmark for the transition between a southern/central lowlands settlement tradition and that of the northern lowlands, with settlement in the southern Chenes region showing attributes of both.

Conclusion

Twentieth-century researchers were indeed correct that interior Campeche is a substantially anthropogenic landscape (*sensu* Chase & Chase 2016). This and other recent analyses (Hernández Gómez 2021; Šprajc *et al.* 2021) make clear that this area was also, in places, an urbanised landscape where rural populations articulated and interacted with crowded cities (Figure 10).

When the regional-scale settlement density values from a vast area of Campeche and western Quintana Roo—55.3strs/km² reported here, 49strs/km² in G-LiHT (Schroder *et al.* 2020), minimally 61strs/km² in the Chactun region (see Šprajc *et al.* 2021)—are considered alongside the discovery of new urban areas such as Valeriana within pseudorandom survey blocks, we can only conclude that cities and dense settlement are simply ubiquitous across large swaths of the central Maya Lowlands. In this regard, the Alianza data are relevant because of the degree to which they do *not* reveal a new and substantially different perspective




Figure 10. Sites and settlement densities in the Alianza survey area (figure by authors).

on Maya settlement density and urbanism when compared with other datasets. Rather, they offer an unambiguous rejoinder to the concern (voiced, for example, by Ringle *et al.* 2021) that archaeologically motivated lidar surveys have inflated estimates of regional settlement density. To the contrary, pseudorandom surveys—those combining expedient and random sampling—like this one, have established that anyone who is waiting for a sparsely settled

Maya hinterland large enough to offset the high settlement densities documented by archaeological lidar surveys is running out of places to look.

As archaeologists work to document and characterise ancient settlement systems throughout the tropics, local scale ‘dense pasts’ are proliferating. It is increasingly clear that dense palimpsests of human modification are not exceptional in these settings, despite long-standing biases that presumed them to be. The field of archaeology now confronts the twin tasks of understanding how these settlement palimpsests accrued through time, and of charting how they varied across space—the better to appreciate just how crowded tropical antiquity may have been.

Data availability statement

Point clouds (.laz), digital elevation models (.tif) and vector points representing our interpretation of archaeological buildings (.kmz) are available through Tulane University’s Lab Archives platform with the permission of Mexico’s Instituto Nacional de Antropología e Historia. They may also be downloaded from the *Ancient Maya Settlement* blog (ancientmayasettlement.com).

Acknowledgements

The authors wish to thank Scott Hutson, Whittaker Schroder and two anonymous reviewers for providing detailed and helpful feedback that significantly improved this manuscript during its journey to publication.

Funding statement

Support for this research was provided by Tulane University Middle American Research Institute through a grant for the MARI-GISLAB by the Hitz Foundation.

References

- ARNAULD, M.C., D. MICHELET & P. NONDÉDÉO. 2013. Living together in Río Bec houses: coresidence, rank, and alliance: noble farmers and weak kings in the Classic Maya Lowlands: the Río Bec Archaeological Project, 2002–2010. *Ancient Mesoamerica* 24: 469–93.
<https://doi.org/10.1017/S0956536114000029>
- AULD-THOMAS, L. 2022. A recipe for simple red relief. *Ancient Maya settlement: investigating how the ancient Maya disposed themselves over the landscape* [Online]. Available at: <https://ancientmayasettlement.com/2022/02/12/a-recipe-for-simple-red-relief/> (accessed 12 February 2022).
- BANNING, E.B. 2021. Sampled to death? The rise and fall of probability sampling in archaeology. *American Antiquity* 86: 43–60.
<https://doi.org/10.1017/aaq.2020.39>
- BARNES, E.B. *et al.* 2009. Las ruinas de Oxpehul, Campeche: su corte real y su huella urbana dentro de la Cuenca de Calakmul, in edited volume *Los investigadores de la cultura Maya*: 194–219. Campeche: Universidad Autónoma de Campeche.
- CANUTO, M.A. & L. AULD-THOMAS. 2020. Reconocimiento y esfuerzos de verificación de lidar en La Corona y El Achiotal, in T. Barrientos Q., M.A. Canuto & M. López (ed.) *Informe Final, Temporada 2019, Proyecto Regional Arqueológico La Corona*: 263–87. New Orleans and Guatemala City: Tulane University and Universidad del Valle, Guatemala.
- 2021. Taking the high ground: a model for lowland Maya settlement patterns. *Journal of Anthropological Archaeology* 64.
<https://doi.org/10.1016/j.jaa.2021.101349>

- 2024. Greater than the sum of its surveyed parts: reconstructing the population of Late Classic Peten with updated methods and new data, in A.S.Z. Chase, A.F. Chase & D.Z. Chase (ed.) *Ancient Mesoamerican population history: demography, social complexity, and change*. 215–66. Tucson: University of Arizona Press.
- CANUTO, M.A. *et al.* 2018. Ancient lowland Maya complexity as revealed by airborne laser scanning of northern Guatemala. *Science* 361. <https://doi.org/10.1126/science.aau0137>
- CHASE, A.F. & D.Z. CHASE. 2016. Urbanism and anthropogenic landscapes. *Annual Review of Anthropology* 45: 361–76. <https://doi.org/10.1146/annurev-anthro-102215-095852>
- CHASE, A.F. *et al.* 2011. Airborne lidar, archaeology, and the ancient Maya landscape at Caracol, Belize. *Journal of Archaeological Science* 38: 387–98. <https://doi.org/10.1016/j.jas.2010.09.018>
- COOK, B. *et al.* 2013. NASA Goddard's LIDAR, hyperspectral and thermal (G-LiHT) airborne imager. *Remote Sensing* 5: 4045–66. <https://doi.org/10.3390/rs5084045>
- DUNNING, N.P. 1992. *Lords of the hills: ancient Maya settlement in the Puuc Region, Yucatan, Mexico* (Monographs in World Archaeology). Madison (WI): Prehistory.
- DUNNING, N.P., T. BEACH, P. FARRELL & S. LUZZADDER-BEACH. 1998. Prehispanic agrosystems and adaptive regions in the Maya lowlands. *Culture and Agriculture*: 87–100. <https://doi.org/10.1525/cag.1998.20.2-3.87>
- DUNNING, N.P. *et al.* 2019. Margin for error: anthropogenic geomorphology of Bajo edges in the Maya lowlands. *Geomorphology* 331: 127–45. <https://doi.org/10.1016/j.geomorph.2018.09.002>
- DUNNING, N.P. & J.K. KOWALSKI. 1994. Lords of the hills: Classic Maya settlement patterns and political iconography in the puuc region, Mexico. *Ancient Mesoamerica* 5: 63–95. <https://doi.org/10.1017/S095653610000105X>
- EATON, J.D. 1975. Ancient agricultural farmsteads in the Río Bec region of Yucatan. *Contributions of the University of California Archaeological Research Facility* 27: 56–82.
- EVANS, D.H. *et al.* 2013. Uncovering archaeological landscapes at Angkor using lidar. *Proceedings of the National Academy of Sciences USA* 110: 12595–600. <https://doi.org/10.1073/pnas.1306539110>
- EVANS, C., O. ALDRED & A. COOPER. 2023. Dense pasts: settlement archaeology after Fox's *The archaeology of the Cambridge region* (1923). *Antiquity* 97: 1193–211. <https://doi.org/10.15184/aqy.2023.99>
- FERNANDEZ-DIAZ, J.C., W.E. CARTER, R.L. SHRESTHA & C.L. GLENNIE. 2014. Now you see it... now you don't: understanding airborne mapping LiDAR collection and data product generation for archaeological research in Mesoamerica. *Remote Sensing* 6: 9951–10001. <https://doi.org/10.3390/rs6109951>
- FISHER, C.T., A.S. COHEN, J.C. FERNÁNDEZ-DÍAZ & S.J. LEISZ. 2017. The application of airborne mapping LiDAR for the documentation of ancient cities and regions in tropical regions. *Quaternary International* 448: 129–38. <https://doi.org/10.1016/j.quaint.2016.08.050>
- FLETCHER, R. 1995. *The limits of settlement growth: a theoretical outline*. Cambridge: Cambridge University Press.
- 2009. Low density, agrarian-based urbanism: a comparative view. *Insights* 2: 1–19.
- 2012. Low-density, agrarian-based urbanism: scale, power and ecology, in M.E. Smith (ed.) *The comparative archaeology of complex societies*: 285–320. New York: Cambridge University Press.
- FREIDEL, D.A., A.F. CHASE, A.S. DOWD & J. MURDOCK. (ed.) 2017. *Maya E groups: calendars, astronomy, and urbanism in the Early Lowlands*. Gainesville: University Press of Florida.
- GARCÍA SÁNCHEZ, J. 2018. Archaeological LiDAR in Italy: enhancing research with publicly accessible data. *Antiquity* 92. <https://doi.org/10.15184/aqy.2018.147>
- GARRISON, T.G., A.E. THOMPSON, S. KRAUSE, S. ESHLEMAN, J.C. FERNANDEZ-DIAZ, J.D. BALDWIN & R. CAMBRANES. 2022. Assessing the lidar revolution in the Maya lowlands: a geographic approach to understanding feature classification accuracy. *Progress in Physical Geography: Earth and Environment* 47. <https://doi.org/10.1177/03091333221138050>
- GOLDEN, C. *et al.* 2016. Reanalyzing environmental lidar data for archaeology: Mesoamerican applications and implications. *Journal of Archaeological Science: Reports* 9: 293–308. <https://doi.org/10.1016/j.jasrep.2016.07.029>
- GUERNSEY, J. 2010. A consideration of the quatrefoil motif in Preclassic Mesoamerica. *RES*:

- Anthropology and Aesthetics* 57/58: 75–96.
<https://doi.org/10.1086/RESvn1ms25769973>
- HERNÁNDEZ GÓMEZ, Q. 2021. La visión LiDAR, el renacimiento de las ciudades en Tierras Bajas Mayas de Campeche: el patrón de asentamiento de El Palmar. Unpublished Bachelors dissertation, Escuela Nacioanl de Antropología e Historia.
- HUTSON, S.R. 2016. *The ancient urban Maya: neighborhoods, inequality, and built form*. Gainesville: University Press of Florida.
- HUTSON, S.R., N.P. DUNNING, B. COOK, T. RUHL, N.C. BARTH & D. CONLEY. 2021. Ancient Maya rural settlement patterns, household cooperation, and regional subsistence interdependency in the Río Bec area: contributions from G-LiHT. *Journal of Anthropological Research* 77: 550–79.
<https://doi.org/10.1086/716750>
- Instituto Nacional de Antropología e Historia. 2020. Con el empleo de tecnología LiDAR y prospección en campo, se registran áreas con monumentos arqueológicos en el municipio de Candelaria, Campeche. *Boletín*: 1–2. Press release available at:
<https://www.gob.mx/cultura/prensa/con-el-empleo-de-tecnologia-lidar-y-prospeccion-en-campo-se-registran-areas-con-monumentos-arqueologicos-en-el-municipio-de-candelaria-campeche> (accessed 12 February 2022).
- JACKMOND, G., D. FONOTI & M.M. TAUTUNU. 2018. Sāmoa's hidden past: lidar confirms inland settlement and suggests larger populations in pre-contact Sāmoa. *Journal of the Polynesian Society* 127: 73–90.
<https://doi.org/10.15286/jps.127.1.73-90>
- KEPECS, S. & S. BOUCHER. 1996. The pre-hispanic cultivation of Rejolladas and Stone-Lands: new evidence from northeast Yucatán, in S.L. Fedick (ed.) *The man-aged mosaic: ancient Maya agriculture and resource use*: 69–91. Salt Lake City: University of Utah Press.
- KOKALJ, Ž. & J. MAST. 2021. Space lidar for archaeology? Reanalyzing GEDI data for detection of ancient Maya buildings. *Journal of Archaeological Science: Reports* 36.
<https://doi.org/10.1016/j.jasrep.2021.102811>
- LEMONNIER, E. & B. VANNIÈRE. 2013. Agrarian features, farmsteads, and homesteads in the Río Bec nuclear zone, Mexico. *Ancient Mesoamerica* 24: 397–413.
<https://doi.org/10.1017/S0956536113000242>
- MAY HAU, J., R. COHUAH MUÑOZ, R. GONZALEZ HEREDIA & W.J. FOLAN. 1990. *The map of the ruins of Calakmul, Campeche, Mexico*. Campeche: Universidad Autónoma de Campeche.
- MERK, S. 2004. Some Maya sites in the northern Río Bec Region, Campeche. *Mexicon* 26: 46–48.
- MERWIN, R.E. 1913. The ruins of the southern part of the peninsula of Yucatan, with special reference to their place in the Maya culture. Unpublished PhD dissertation, Harvard University.
- PÉREZ RIVAS, M.E., A.L. CORRAL, Á.J.G. UITZ & A.V. MORLET. 2023. Un sitio monumental en Campeche. *Exploraciones en el sitio S.C.R. Arqueología Mexicana* edición especial: 84–89.
- PERIPATO, V. *et al.* 2023. More than 10,000 pre-Columbian earthworks are still hidden throughout Amazonia. *Science* 382: 103–9.
<https://doi.org/10.1126/science.ade2541>
- PLOG, S. & K.V. FLANNERY. 1976. Sampling on the regional level, in K.V. Flannery (ed.) *The early Mesoamerican village*: 131–60. New York: Academic Press.
- PRÜMERS, H., C.J. BETANCOURT, J. IRIARTE, M. ROBINSON & M. SCHAICH. 2022. Lidar reveals pre-Hispanic low-density urbanism in the Bolivian Amazon. *Nature* 606: 325–8.
<https://doi.org/10.1038/s41586-022-04780-4>
- REESE-TAYLOR, K. *et al.* 2016. Boots on the ground at Yaxnohc'ah: ground-truthing LiDAR in a complex tropical landscape. *Advances in Archaeological Practice* 4: 314–38.
<https://doi.org/10.7183/2326-3768.4.3.314>
- RICE, D.S. & D.E. PULESTON. 1981. Ancient Maya settlement patterns in the Peten, Guatemala, in W.A. Ashmore (ed.) *Lowland Maya settlement patterns*: 121–56. Albuquerque: University of New Mexico Press.
- RINGLE, W.M., T.G. NEGRÓN, R.M. CIAU, K.E. SELIGSON, J.C. FERNANDEZ-DIAZ & D. ORTEGÓN ZAPATA. 2021. Lidar survey of ancient Maya settlement in the Puuc region of Yucatan, Mexico. *PLoS ONE* 16.
<https://doi.org/10.1371/journal.pone.0249314>
- ROSTAIN, S. *et al.* 2024. Two thousand years of garden urbanism in the Upper Amazon. *Science* 383: 183–9.
<https://doi.org/10.1126/science.adi6317>
- RUPPERT, K.J. & J.H. DENISON. 1943. *Archaeological reconnaissance in Campeche, Quintana Roo, and Peten*. Washington (D.C.): Carnegie Institute of Washington.

- SCHRODER, W. *et al.* 2020. The lowland Maya settlement landscape: environmental LiDAR and ecology. *Journal of Archaeological Science: Reports* 33. <https://doi.org/10.1016/j.jasrep.2020.102543>
- SELIGSON, K.E., S. ORTIZ RUIZ & L. BARBA PINGARRÓN. 2019. Prehispanic Maya burnt lime industries: previous studies and future directions. *Ancient Mesoamerica* 30: 199–219. <https://doi.org/10.1017/S0956536117000347>
- ŠPRAJC, I. *et al.* 2021. Ancient Maya water management, agriculture, and society in the area of Chactún, Campeche, Mexico. *Journal of Anthropological Archaeology* 61. <https://doi.org/10.1016/j.jaa.2020.101261>
- STAUFFER, J.G., S.B. GROOMS, L.W. HU, J. MERSMANN, T.R. KIDDER & E.R. HENRY. 2023. Reimagining the development of downtown Cahokia using remote sensing visualizations from the western edge of the Grand Plaza. *Land* 12. <https://doi.org/10.3390/land12020342>
- SUÁREZ AGUILAR, V. 2007. *Rescates y salvamentos arqueológicos en Campeche y Yucatán*. Mexico: CONACULTA–INAH.
- THOMAS, P.M. 1981. *Prehistoric Maya settlement: patterns at Becan, Campeche, Mexico*. New Orleans (LA): Middle American Research Institute, Tulane University.
- THOMPSON, A.E. 2020. Detecting Classic Maya settlements with lidar-derived relief visualizations. *Remote Sensing* 12. <https://doi.org/10.3390/rs12172838>
- THOMPSON, V.D. 2023. Considering urbanism at Mound Key (*Caalus*), the capital of the Calusa in the 16th century, southwest Florida, USA. *Journal of Anthropological Archaeology* 72. <https://doi.org/10.1016/j.jaa.2023.101546>
- TURNER II, B.L. 1978. Ancient agricultural land use in the central lowlands, in P.D. Harrison & B.L. Turner II (ed.) *Pre-Hispanic Maya agriculture*: 163–83. Albuquerque: University of New Mexico Press.
- 1983. *Once beneath the forest: prehistoric terracing in the Río Bec Region of the Maya Lowlands*. London: Routledge.
- VERBOVSEK, T., T. POPIT & Ž. KOKALJ. 2019. VAT method for visualization of mass movement features: an alternative to hillshaded DEM. *Remote Sensing* 11. <https://doi.org/10.3390/rs11242946>
- WALKER, W.S. 2015. La iniciativa LiDAR aerotransportado de la Alianza México REDD+. Mexico: The Nature Conservancy, México. Available at: <https://www.tncmx.org/content/dam/tnc/nature/en/documents/mexico/M-REDD-FichaIniciativaLiDARREDD.pdf> (accessed 12 February 2022).
- WEBSTER, D.L. 2018. *The population of Tikal: implications for Maya demography* (Paris Monographs in American Archaeology). Oxford: Archaeopress.
- WITSCHHEY, W.R.T. & C.T. BROWN. 2014. 5,000 sites and counting: the inspiration of Maya settlement studies, in G.E. Braswell (ed.) *The ancient Maya of Mexico: reinterpreting the past of the northern Maya Lowlands*: 184–202. London: Routledge.